Amendments to the Claims

This listing of claims will replace all prior versions, and listings, of claims in the application:

Listing of Claims:

1 1. (currently amended): A system for diagnosing and monitoring 2 myocardial ischemia for automated remote patient care, comprising: 3 a database storing a plurality of monitoring sets which each comprise recorded measures relating to patient information recorded on a substantially 4 5 continuous basis: б a server retrieving and processing a plurality of the monitoring sets, 7 comprising: 8 a comparison module determining a patient status change by 9 comparing at least one recorded measure and time from each of the monitoring 10 sets to at least one other recorded measure and time from another monitoring set 11 with both recorded measures relating to a same type of patient information 12. recorded at different times; and 13 an analysis module testing each comparing the patient status 14 change for an absence, an onset, a progression, a regression, and a status que of 15 myocardial ischemia against to a predetermined indicator threshold, including at 16 least one hysteresis threshold, corresponding to same type of patient information 17 as the recorded measures which were compared and to at least one stickiness 18 threshold, the indicator threshold corresponding to a quantifiable physiological 19 measure of a pathophysiology indicative of myocardial ischemia and the 20 hysteresis stickiness threshold corresponding to a diagnosis or treatment based on 21 a duration and degree of the patient status change and determining an onset, a 22 progression, a regression, and a status quo of myocardial ischemia dependent 23 upon the comparison.

> 2. (original): A system according to Claim 1, further comprising:

OA Response

1

2	an analysis submodule managing the myocardial ischemia and outcomes
3	thereof through administering at least one of anticoagulation, antiplatelet drugs,
4	beta-blockade, coronary vasodilators, afterload reduction, lipid lowering drugs,
5	electrical therapies, and mechanical therapies.
1	 (original): A system according to Claim 1, further comprising:
2	a database submodule periodically receiving a monitoring set comprising
3	recorded measures for an individual patient, each recorded measure in the
4	monitoring set having been recorded by at least one of a medical device adapted
5	to be implanted in an individual patient and an external medical device proximal
6	to the individual patient when the device measures are recorded and storing the
7	received monitoring set in the database as part of a patient care record for the
8	individual patient.
1	4. (original): A system according to Claim 3, further comprising:
2	a set of further indicator thresholds, each indicator threshold
3	corresponding to a quantifiable physiological measure used to detect a
4	pathophysiology indicative of diseases other than myocardial ischemia;
5	a comparison submodule comparing each patient status change to each
б	such further indicator threshold corresponding to the same type of patient
7	information as the at least one recorded measure and the at least one other
8	recorded measure; and
9	an analysis submodule testing each patient status change against each such
10	further indicator threshold corresponding to the same type of patient information
11	as the recorded measures which were compared
1	5. (original): A system according to Claim 1, further comprising:
2	a comparison submodule determining a change in patient status by
3	comparing at least one recorded quality of life measure to at least one other

4

corresponding recorded quality of life measure.

1

3 4

> 1 2

> 3

4

5

1

2 3

6.	(currently amended): A system according to Claim 1, further
comprising: w	herein the stickiness threshold comprises a set of stickiness
indicators for	each type of patient information, each stickiness indicator
corresponding	to a temporal limit related to a program of patient diagnosis or
treatment[[;]]	further comprising:
a comp	parison submodule comparing a time span occurring between each
patient status	change for each recorded measure to the stickiness indicator relating
to the same typ	pe of patient information as the recorded measure being compared;
and ·	•
an ana	lysis submodule determining a revised program of patient diagnosis
or treatment re	esponsive to each patient status change occurring subsequent to a
time span exc	eeding the stickiness indicator.

- 7. (original): A system according to Claim 1, further comprising: a database module retrieving the plurality of monitoring sets from one of a 2 patient care record for an individual patient, a peer group, and a overall patient population.
 - (original): A system according to Claim 1, further comprising: 8. the database further storing a reference baseline comprising recorded measures which each relate to patient information recorded during an initial time period and comprise either medical device measures or derived measures a database submodule obtaining at least one of the reference baseline.
 - 9. (original): A system according to Claim 1, wherein the indicator thresholds relate to at least one of a finding of reduced exercise capacity, respiratory distress and angina.
- 10. (original): A system according to Claim 9, wherein the indicator 1 2 thresholds relating to the finding of reduced exercise capacity are selected from the group consisting of decreased cardiac output, decreased mixed venous oxygen 3 4 score, and decreased patient activity score.

(original): A system according to Claim 9, wherein the indicator

1

11.

	· • · · · ·
2	thresholds relating to the finding of respiratory distress are selected from the
3	group consisting of increased respiratory rate, decreased transthoracic impedance,
4	and increased pulmonary artery diastolic pressure.
5	12. (original): A system according to Claim 9, wherein the indicator
б	thresholds relating to the finding of angina are selected from the group consisting
7	of increased ST segment, decreased ST segment, increased myocardial creatinine
8	kinase, increased troponin, increased coronary sinus lactate, changed myocardial
9	wall motion, ventricular tachycardia, T wave changes, and QRS changes.
1	13. (currently amended): A method for diagnosing and monitoring
2	myocardial ischemia for automated remote patient care, comprising:
3	storing a plurality of monitoring sets which each comprise recorded
4	measures relating to patient information recorded on a substantially continuous
5	basis in a database;
6	retrieving a plurality of the monitoring sets from the database;
7	determining a patient status change by comparing at least one recorded
8	measure and time from each of the monitoring sets to at least one other recorded
9	measure and time from another monitoring set with both recorded measures
10	relating to a same type of patient information recorded at different times; and
11	testing each comparing the patient status change for an absence, an onset,
12	a progression, a regression, and a status quo of myocardial ischemia against to a
13	predetermined indicator threshold, including at least one hysteresis threshold,
14	corresponding to a same type of patient information as the recorded measures
15	which were compared and to at least one stickiness threshold, the indicator
16	threshold corresponding to a quantifiable physiological measure of a
17	pathophysiology indicative of myocardial ischemia and the hysteresis stickiness
18	threshold corresponding to a diagnosis or treatment based on a duration and
19	degree of the patient status change and determining an onset, a propression, a

20	regression, and a status quo of myocardial ischemia dependent upon the
21	comparison.
1	14. (original): A method according to Claim 13, further comprising:
2	managing the myocardial ischemia and outcomes thereof through
3	administration of at least one of anticoagulation, antiplatelet drugs, beta-blockade
4	coronary vasodilators, afterload reduction, lipid lowering drugs, electrical
5	therapies, and mechanical therapies.
1	15. (original): A method according to Claim 13, further comprising:
2	periodically receiving a monitoring set for an individual patient, each
3	recorded measure in the monitoring set having been recorded by at least one of a
4	medical device adapted to be implanted in an individual patient and an external
5	medical device proximal to the individual patient when the device measures are
6	recorded; and
7	storing the received monitoring set in the database as part of a patient care
.8	record for the individual patient.
1	16. (original): A method according to Claim 15, further comprising:
2	defining a set of further indicator thresholds, each indicator threshold
3	corresponding to a quantifiable physiological measure used to detect a
4	pathophysiology indicative of diseases other than myocardial ischemia;
5	comparing each patient status change to each such further indicator
6	threshold corresponding to the same type of patient information as the at least on
7	recorded measure and the at least one other recorded measure; and
8	testing each patient status change against each such further indicator
9	threshold corresponding to the same type of patient information as the recorded
10	measures which were compared.
1	17. (original): A method according to Claim 13, further comprising:

2	determining a change in patient status by comparing at least one recorded
3	quality of life measure to at least one other corresponding recorded quality of life
4	measure.
1	18. (currently amended): A method according to Claim 13, further
2	comprising: defining wherein the stickiness threshold comprises a set of stickines
3	indicators for each type of patient information, each stickiness indicator
4	corresponding to a temporal limit related to a program of patient diagnosis or
5	treatment[[+]] further comprising:
6	comparing a time span occurring between each patient status change for
7	each recorded measure to the stickiness indicator relating to the same type of
8	patient information as the recorded measure being compared; and
9	determining a revised program of patient diagnosis or treatment
10	responsive to each patient status change occurring subsequent to a time span
11	exceeding the stickiness indicator.
1	19. (original): A method according to Claim 13, further comprising:
2	retrieving the plurality of monitoring sets from one of a patient care record
3	for an individual patient, a peer group, and a overall patient population.
1	20. (original): A method according to Claim 13, further comprising:
2	retrieving a reference baseline comprising recorded measures which each
3	relate to patient information recorded during an initial time period and comprise
4	either medical device measures or derived measures calculable therefrom; and
5	obtaining at least one of the at least one recorded measure and the at least
6	one other recorded measure from the reference baseline.
1	21. (currently amended): An automated A method according to Claim
2	13, wherein the indicator thresholds relate to at least one of a finding of reduced
3	exercise capacity, respiratory distress and angina.
4	22. (original): A method according to Claim 21, wherein the indicator
5	thresholds relating to the finding of reduced exercise capacity are selected from

3

4

5

6

7

8

9

10

11

12

13

14

15

16

17

18

Response to First Office Action Docket No. 020.0340.US.CON

- the group consisting of decreased cardiac output, decreased mixed venous oxygen
 score, and decreased patient activity score.
- 23. (original): A method according to Claim 21, wherein the indicator thresholds relating to the finding of respiratory distress are selected from the group consisting of increased respiratory rate, decreased transthoracic impedance, and increased pulmonary artery diastolic pressure.
- 24. (original): A method according to Claim 21, wherein the indicator thresholds relating to the finding of angina are selected from the group consisting of increased ST segment, decreased ST segment, increased myocardial creatinine kinase, increased troponin, increased coronary sinus lactate, changed myocardial wall motion, ventricular tachycardia, T wave changes, and QRS changes.
 - 25. (currently amended): A computer-readable storage medium holding code for diagnosing and monitoring myocardial ischemia for automated remote patient care, comprising:

code for storing a plurality of monitoring sets which each comprise recorded measures relating to patient information recorded on a substantially continuous basis in a database;

code for retrieving a plurality of the monitoring sets from the database;

code for determining a patient status change by comparing at least one
recorded measure and time from each of the monitoring sets to at least one other
recorded measure and time from another monitoring set with both recorded
measures relating to a same type of patient information recorded at different
times; and

code for testing each comparing the patient status change for an absence, an onset, a progression, a regression, and a status quo of myocardial ischemia against to a predetermined indicator threshold, including at least one hysteresis threshold, corresponding to a same type of patient information as the recorded measures which were compared and to at least one stickiness threshold, the indicator threshold corresponding to a quantifiable physiological measure of a

19	pathophysiology indicative of myocardial ischemia and the hysteresis stickiness
20	threshold corresponding to a diagnosis or treatment based on a duration and
21	degree of the patient status change and code for determining an onset, a
22	progression, a regression, and a status quo of myocardial ischemia dependent
23	upon the comparison.
1	26. (original): A storage medium according to Claim 25, further
2	comprising:
3	code for managing the myocardial ischemia and outcomes thereof through
4	administration of at least one of anticoagulation, antiplatelet drugs, beta-blockade,
5	coronary vasodilators, afterload reduction, lipid lowering drugs, electrical
6	therapies, and mechanical therapies.
v	therapies, and mechanical dictapies.
1	27. (original): A storage medium according to Claim 25, further
2	comprising:
3	code for periodically receiving a monitoring set for an individual patient,
4	each recorded measure in the monitoring set having been recorded by at least one
5	of a medical device adapted to be implanted in an individual patient and an
б	external medical device proximal to the individual patient when the device
7	measures are recorded; and
8	code for storing the received monitoring set in the database as part of a
9	patient care record for the individual patient.
1	28. (original): A storage medium according to Claim 27, further
2	comprising:
3	code for defining a set of further indicator thresholds, each indicator
4	threshold corresponding to a quantifiable physiological measure used to detect a
5	pathophysiology indicative of diseases other than myocardial ischemia;
6	code for comparing each patient status change to each such further
7	
8	indicator threshold corresponding to the same type of patient information as the at
Ü	least one recorded measure and the at least one other recorded measure; and

9	code for testing each patient status change against each such further
10	indicator threshold corresponding to the same type of patient information as the
11	recorded measures which were compared
1	29. (original): A storage medium according to Claim 25, further
2	comprising:
3	code for determining a change in patient status by comparing at least one
4	recorded quality of life measure to at least one other corresponding recorded
5	quality of life measure.
1	30. (currently amended): A storage medium according to Claim 25,
2	further comprising: code for defining wherein the stickiness threshold comprises
3	set of stickiness indicators for each type of patient information, each stickiness
4	indicator corresponding to a temporal limit related to a program of patient
5	diagnosis or treatment[[i]] further comprising:
6	code for comparing a time span occurring between each patient status
.7	change for each recorded measure to the stickiness indicator relating to the same
8	type of patient information as the recorded measure being compared; and
9	code for determining a revised program of patient diagnosis or treatment
10	responsive to each patient status change occurring subsequent to a time span
. 11	exceeding the stickiness indicator.
1	31. (original): A storage medium according to Claim 25, further
2	comprising:
3	code for retrieving the plurality of monitoring sets from one of a patient
4	care record for an individual patient, a peer group, and a overall patient
5	population.
1	32. (original): A storage medium according to Claim 25, further
2	comprising:
3	code for retrieving a reference baseline comprising recorded measures
4	Which each relate to nations information recorded during an initial time period on

5	comprise either medical device measures or derived measures calculable
6	therefrom; and
7	code for obtaining at least one of the at least one recorded measure and the
8	at least one other recorded measure from the reference baseline.
9	33. (currently amended): An automated collection and analysis patient
10	care system for diagnosing and monitoring myocardial ischemia and outcomes
11	thereof, comprising:
12	a database storing patient monitoring information, comprising:
13	a plurality of monitoring sets, each monitoring set comprising
14	recorded measures which each relate to patient information and comprise either
15	medical device measures or derived measures calculable therefrom, the medical
16	device measures having been regularly recorded on a substantially continuous
17	basis;
18	a set of stored indicator thresholds, each indicator threshold
19	corresponding to a quantifiable physiological measure of a pathophysiology
20	indicative of myocardial ischemia and relating to a same type of patient
21	information as at least one of the recorded measures;
22	a server diagnosing a myocardial ischemia finding comprising an absence,
23	an onset, a progression, a regression, and a status quo of myocardial ischemia,
24	comprising:
25	a receiver retrieving a plurality of the monitoring sets from the
26	database;
27	an analysis module determining a change in patient status by
28	comparing at least one recorded measure and time to at least one other recorded
29	measure and time from another monitoring set with both recorded measures
30	relating to a same type of patient information recorded at different times; and
31	a comparison module comparing each patient status change to the
32	indicator threshold, including at least one hysteresis threshold, corresponding to a
33	same type of patient information as the recorded measures which were compared

4

5

1

2

3

4

5

6

7

Response to First Office Action Docket No. 020.0340.US.CON

- and the hysteresis to at least one stickiness threshold corresponding to a diagnosis or treatment based on a duration and degree of the patient status change.
- 1 34. (original): A system according to Claim 33, wherein the device
 2 measures are recorded by at least one of a medical device adapted to be implanted
 3 in an individual patient and an external medical device proximal to the individual
 4 patient when the device measures are recorded.
- 35. (original): A system according to Claim 33, wherein each of the
 monitoring sets comprises recorded measures relating to patient information
 solely for the individual patient, further comprising:
 - a database module retrieving each monitoring set from a patient care record for the individual patient and obtaining the at least one recorded measure and the at least one other recorded measure from the retrieved monitoring sets.
 - 36. (original): A system according to Claim 33, wherein each of the monitoring sets comprises recorded measures relating to patient information for a peer group of patients to which the individual patient belongs, further comprising: a database module retrieving at least one monitoring set from a patient

care record for the individual patient, retrieving at least one other monitoring set

- from a patient care record in the same patient peer group, and obtaining the at least one recorded measure from the at least one monitoring set and the at least
- 8 one other recorded measure from the at least one other monitoring set.
- 1 37. (original): A system according to Claim 33, wherein each of the 2 monitoring sets comprises recorded measures relating to patient information for a 3 general population of patients, further comprising:
- a database module retrieving at least one monitoring set from a patient
- 5 care record for the individual patient, retrieving at least one other monitoring set
- 6 from a patient care record in the overall patient population, and obtaining the at
- 7 least one recorded measure from the at least one monitoring set and the at least
- 8 one other recorded measure from the at least one other monitoring set.

1	38. (original): A system according to Claim 33, further comprising:
2	a database submodule further storing a reference baseline comprising
3	recorded measures which each relate to patient information recorded by the
4	medical device adapted to be implanted during an initial time period and comprise
5	cither device measures recorded by the medical device adapted to be implanted or
6	derived measures calculable therefrom; and
7	a database submodule obtaining at least one of the at least one recorded
8	measure and the at least one other recorded measure from the reference baseline.
1	39. (original): A system according to Claim 38, wherein the reference
2	baseline comprises recorded measures relating to patient information for one of
3	the individual patients solely, a peer group of patients to which the individual
4	patient belongs, and a general population of patients.
1	40. (original): A system according to Claim 33, wherein the indicator
2	thresholds relate to a priori limits selected from the group comprising ST segment
3	elevation, myocardial band creatinine kinase mass, and troponin levels.
1	41. (original): A system according to Claim 40, wherein the ST
2	segment elevation exceeds substantially 2.0 mm in an absence of a QRS duration
3	greater than or equal to substantially 120 ms, the myocardial band creatinine
4	kinase mass exceeds substantially 5 ng/ml, and the troponin-I exceeds
5	substantially 0.5 ng/ml.
1	42. (original): A system according to Claim 33, the comparison
2	module further comprising:
3	a comparison submodule grading the comparisons between each patient
4	status change and corresponding indicator threshold on a fixed scale based on a
5	degree of deviation from the indicator threshold; and
6	a comparison submodule determining an overall patient status change by
7	performing a summation over the individual graded comparisons.

1	43. (original): A system according to Claim 33, the comparison
2	module further comprising:
3	a module determining probabilistic weightings of the comparisons
4	between each patient status change and corresponding indicator threshold based
5	on a statistical deviation and trends via linear fits from the indicator threshold;
6	and .
7	a comparison submodule determining an overall patient status change by
8	performing a summation over the individual graded comparisons.
1	44. (original): A system according to Claim 33, wherein each
2	monitoring set further comprises quality of life and symptom measures recorded
3	by the individual patient, the server further comprising:
4	a quality of life module determining a change in patient status by
5	comparing at least one recorded quality of life measure to at least one other
6	corresponding recorded quality of life measure; and
7	the server incorporating each patient status change in quality of life into
8	the myocardial ischemia finding to either refute or support the diagnosis.
1	45. (original): A system according to Claim 33, further comprising:
2	a set of stored further indicator thresholds, each indicator threshold
3	corresponding to a quantifiable physiological measure used to detect a
4	pathophysiology indicative of diseases other than myocardial ischemia of disease;
5	and
6	a diagnostic submodule diagnosing a finding of a disease other than
7	myocardial ischemia, the comparison module further comprising comparing each
8	patient status change to each such further indicator threshold corresponding to the
9	same type of patient information as the at least one recorded measure and the at
10	least one other recorded measure.
1	46. (currently amended): A system according to Claim 33, further
2	comprising: wherein the stickiness threshold comprises a set of stickiness

3	indicators, each indicator threshold corresponding to a temporal limit related to a
4	course of patient care[[; and]] further comprising:
5	a feedback module comparing a time span between each patient status
6	change for each recorded measure to the stickiness indicator corresponding to the
7	same type of patient information as the recorded measure being compared.
1	47. (original): A system according to Claim 33, further comprising:
2	a feedback module providing automated feedback to the individual patient
3	when a myocardlal ischemia finding is indicated.
1	48. (original): A system according to Claim 47, further comprising:
2	the feedback module performing an interactive dialogue between the
3	individual patient and the patient care system regarding a medical condition of the
4	individual patient.
1	49. (currently amended): A method for diagnosing and monitoring
2	myocardial ischemia and outcomes using an automated collection and analysis
3	patient care system, comprising:
4	storing a plurality of monitoring sets in a database, each monitoring set
5	comprising recorded measures which each relate to patient information and
б	comprise either medical device measures or derived measures calculable
7	therefrom, the medical device measures having been regularly recorded on a
8	substantially continuous basis;
9	retrieving a plurality of the monitoring sets from the database;
10	defining a set of indicator thresholds, each indicator threshold
11	corresponding to a quantifiable physiological measure of a pathophysiology
12 ·	indicative of myocardial ischemia and relating to a same type of patient
13	information as at least one of the recorded measures; and
14	diagnosing a myocardial ischemia finding comprising an absence, an
15	onset, a progression, a regression, and a status quo of myocardial ischemia

comprising:

. 16

17	determining a change in patient status by comparing at least one
18	recorded measure and time to at least one other recorded measure and time from
19	another monitoring set with both recorded measures relating to a same type of
20	patient information recorded at different times; and
21	comparing each patient status change to the indicator threshold;
22	including at least one hysteresis threshold, corresponding to a same type of patien
23	information as the recorded measures which were compared and the hysteresis to
24	at least one stickiness threshold corresponding to a diagnosis or treatment based
25	on a duration and degree of the patient status change.
1	50. (original): A method according to Claim 49, wherein the device
2	measures are recorded by at least one of a medical device adapted to be implanted
3	in an individual patient and an external medical device proximal to the individual
4	patient when the device measures are recorded.
1	51. (original): A method according to Claim 49, wherein each of the
2	monitoring sets comprises recorded measures relating to patient information
3	solely for the individual patient, further comprising:
4	retrieving each monitoring set from a patient care record for the individual
5	patient; and
6	obtaining the at least one recorded measure and the at least one other
7	recorded measure from the retrieved monitoring sets.
1	52. (original): A method according to Claim 49, wherein each of the
2	monitoring sets comprises recorded measures relating to patient information for a
3	peer group of patients to which the individual patient belongs, further comprising
4	retrieving at least one monitoring set from a patient care record for the
5	individual patient;
6	retrieving at least one other monitoring set from a patient care record in

the same patient peer group; and

8	obtaining the at least one recorded measure from the at least one
9	monitoring set and the at least one other recorded measure from the at least one
10	other monitoring set.
1	53. (original): A method according to Claim 49, wherein each of the
2	monitoring sets comprises recorded measures relating to patient information for a
3	general population of patients, further comprising:
4	retrieving at least one monitoring set from a patient care record for the
5	individual patient;
6	retrieving at least one other monitoring set from a patient care record in
7	the overall patient population; and
8	obtaining the at least one recorded measure from the at least one
9	monitoring set and the at least one other recorded measure from the at least one
10	other monitoring set.
1	54. (original): A method according to Claim 49, further comprising:
2	retrieving a reference baseline comprising recorded measures which each
3	relate to patient information recorded by the medical device adapted to be
4	implanted during an initial time period and comprise either device measures
5	recorded by the medical device adapted to be implanted or derived measures
6	calculable therefrom; and
7	obtaining at least one of the at least one recorded measure and the at least
8	one other recorded measure from the reference baseline.
1	55. (original): A method according to Claim 54, wherein the reference

- baseline comprises recorded measures relating to patient information for one of 3 the individual patients solely, a peer group of patients to which the individual 4 patient belongs, and a general population of patients.
- 1 56. (original): A method according to Claim 49, wherein the indicator 2 thresholds relate to a priori limits selected from the group comprising ST segment 3 elevation, myocardial band creatinine kinase mass, and troponin levels.

2

t	57. (original): A method according to Claim 56, wherein the ST
2	segment elevation exceeds substantially 2.0 mm in an absence of a QRS duration
3	greater than or equal to substantially 120 ms, the myocardial band creatinine
ļ	kinase mass exceeds substantially 5 ng/ml, and the troponin-I exceeds
5	substantially 0.5 ng/ml.
l	58. (original): A method according to Claim 49, the operation of
2	comparing each patient status change further comprising:
3	grading the comparisons between each patient status change and
ļ	corresponding indicator threshold on a fixed scale based on a degree of deviation
5	from the indicator threshold; and
ó	determining an overall patient status change by performing a summation
7	over the individual graded comparisons.
l	59. (original): A method according to Claim 49, the operation of
2	comparing each patient status change further comprising:
3	determining probabilistic weightings of the comparisons between each
Ļ	patient status change and corresponding indicator threshold based on a statistical
5	deviation and trends via linear fits from the indicator threshold; and
5	determining an overall patient status change by performing a summation
7	over the individual graded comparisons.
L	60. (original): A method according to Claim 49, wherein each
?	monitoring set further comprises quality of life and symptom measures recorded
3	by the individual patient, the operation of diagnosing a myocardial ischemia
ļ	finding further comprising:
;	determining a change in patient status by comparing at least one recorded
5	quality of life measure to at least one other corresponding recorded quality of life
,	measure; and
3	incorporating each patient status change in quality of life into the
)	myocardial ischemia finding to either refute or support the diagnosis.

8

1

2

3

4

6

1

2

3

4

5

6

61	. (original): A method according to Claim 49, further comprising:
det	fining a set of further indicator thresholds, each indicator threshold
correspon	ling to a quantifiable physiological measure used to detect a
pathophys	iology indicative of diseases other than myocardial ischemia; and
dia	gnosing a finding of the disease other than myocardial ischemia,
comprising	g comparing each patient status change to each such further indicator
threshold	corresponding to the same type of patient information as the at least one
recorded r	neasure and the at least one other recorded measure.

- 62. (currently amended): A method according to Claim 49, further comprising: defining wherein the stickiness threshold comprises a set of stickiness indicators, each indicator threshold corresponding to a temporal limit related to a course of patient care[[; and]] further comprising:
- 5 comparing a time span between each patient status change for each recorded measure to the stickiness indicator corresponding to the same type of 7 patient information as the recorded measure being compared.
 - 63. (original): A method according to Claim 49, further comprising: providing automated feedback to the individual patient when a myocardial ischemia finding is indicated.
- 1 (original): A method according to Claim 63, further comprising: 2 performing an interactive dialogue between the individual patient and the 3 patient care system regarding a medical condition of the individual patient.
 - 65. (currently amended): A computer-readable storage medium holding code for diagnosing and monitoring myocardial ischemia using an automated collection and analysis patient care system, comprising:
- 7 code for storing a plurality of monitoring sets in a database, each 8 monitoring set comprising recorded measures which each relate to patient 9 information and comprise either medical device measures or derived measures

10	calculable therefrom, the medical device measures having been regularly recorded
11	on a substantially continuous basis;
12	code for retrieving a plurality of the monitoring sets from the database;
13	code for defining a set of indicator thresholds, each indicator threshold
14	corresponding to a quantifiable physiological measure of a pathophysiology
15	indicative of myocardial ischemia and relating to a same type of patient
16	information as at least one of the recorded measures; and
17	code for diagnosing a myocardial ischemia finding comprising an absence
18	an onset, a progression, a regression, and a status quo of myocardial ischemia,
19	comprising:
20	code for determining a change in patient status by comparing at
21	least one recorded measure and time to at least one other recorded measure and
22	time from another monitoring set with both recorded measures relating to a same
23	type of patient information recorded at different times; and
24	code for comparing each patient status change to the indicator
25	threshold, including at least one hysteresis threshold, corresponding to a same
26	type of patient information as the recorded measures which were compared and
27	the hysteresis to at least one stickiness threshold corresponding to a treatment or
28	diagnosis based on a duration and degree of the patient status change.
1	66. (original): A storage medium according to Claim 65, wherein each
2	of the monitoring sets comprises recorded measures relating to patient
3	information solely for the individual patient, further comprising:
4	code for retrieving each monitoring set from a patient care record for the
5	individual patient; and
6	code for obtaining the at least one recorded measure and the at least one
7	other recorded measure from the retrieved monitoring sets.
1	67. (original): A storage medium according to Claim 65, wherein each
2	of the monitoring sets comprises recorded measures relating to natical

3	information for a peer group of patients to which the individual patient octongs,
4	further comprising:
5	code for retrieving at least one monitoring set from a patient care record
6	for the individual patient;
7	code for retrieving at least one other monitoring set from a patient care
8	record in the same patient peer group; and
9	code for obtaining the at least one recorded measure from the at least one
10	monitoring set and the at least one other recorded measure from the at least one
11	other monitoring set.
1	68. (original): A storage medium according to Claim 65, wherein each
2	of the monitoring sets comprises recorded measures relating to patient
3	information for a general population of patients, further comprising:
4	code for retrieving at least one monitoring set from a patient care record
5	for the individual patient;
6	code for retrieving at least one other monitoring set from a patient care
7	record in the overall patient population; and
8	code for obtaining the at least one recorded measure from the at least one
9	monitoring set and the at least one other recorded measure from the at least one
10	other monitoring set.
1	69. (original): A storage medium according to Claim 65, further
2	comprising:
3	code for retrieving a reference baseline comprising recorded measures
4	which each relate to patient information recorded by the medical device adapted
5	to be implanted during an initial time period and comprise either device measures
6	recorded by the medical device adapted to be implanted or derived measures
7	calculable therefrom; and
8	code for obtaining at least one of the at least one recorded measure and the
9	at least one other recorded measure from the reference baseline.

1	70. (original): A storage medium according to Claim 65, the operation
2	of comparing each patient status change further comprising:
3	code for grading the comparisons between each patient status change and
4	corresponding indicator threshold on a fixed scale based on a degree of deviation
5	from the indicator threshold; and
6	code for determining an overall patient status change by performing a
7	summation over the individual graded comparisons.
1	71. (original): A storage medium according to Claim 65, the operation
2	of comparing each patient status change further comprising:
3	code for determining probabilistic weightings of the comparisons between
4	each patient status change and corresponding indicator threshold based on a
5	statistical deviation and trends via linear fits from the indicator threshold; and
6	code for determining an overall patient status change by performing a
7	summation over the individual graded comparisons.
1	72. (original): A storage medium according to Claim 65, wherein each
2	monitoring set further comprises quality of life and symptom measures recorded
3	by the individual patient, the operation of diagnosing a myocardial ischemia
4	finding further comprising:
5	code for determining a change in patient status by comparing at least one
6	recorded quality of life measure to at least one other corresponding recorded
7	quality of life measure; and
8	code for incorporating each patient status change in quality of life into the
9	myocardial ischemia finding to either refute or support the diagnosis.
1	73. (original): A storage medium according to Claim 65, further
2	comprising:
3	code for defining a set of further indicator thresholds, each indicator
4	threshold corresponding to a quantifiable physiological measure used to detect a
5	pathophysiology indicative of diseases other than myocardial ischemia; and

Response to First Office Action Docket No. 020.0340.US.CON

code for diagnosing a finding of the disease other than myocardial
ischemia, comprising comparing each patient status change to each such further
indicator threshold corresponding to the same type of patient information as the at
least one recorded measure and the at least one other recorded measure.
74. (original): A storage medium according to Claim 65, further
comprising: code for defining wherein the stickiness threshold comprises a set of
stickiness indicators, each indicator threshold corresponding to a temporal limit
related to a course of patient care[[; and]] further comprising:
code for comparing a time span between each patient status change for
each recorded measure to the stickiness indicator corresponding to the same type
of patient information as the recorded measure being compared.
75. (original): A storage medium according to Claim 65, further
comprising:
code for providing automated feedback to the individual patient when a
myocardial ischemia finding is indicated.
76. (original): A storage medium according to Claim 75, further
comprising:
code for performing an interactive dialogue between the individual patient
and the patient care system regarding a medical condition of the individual
patient.
77. (currently amended): An automated patient care system for
diagnosing and monitoring myocardial ischemia and outcomes thereof,
comprising:
a medical device regularly recording measures relating to at least one of
monitoring angina, reduced exercise capacity and respiratory distress;
a database maintaining information for an individual patient, comprising

organizing a plurality of monitoring sets in a database, and storing the recorded

8	measures for the individual patient on a substantially continuous basis into a
9	monitoring set in the database;
10	a server evaluating a finding of at least one of an absence, an onset, a
11	progression, a regression, and a status quo of myocardial ischemia, comprising:
12	a comparison module determining a patient status change by
13	comparing at least one recorded measure and time from each of the monitoring
14	sets to at least one other recorded measure and time from another monitoring set
15	with both recorded measures relating to a same type of patient information
16	recorded at different times; and
17	an analysis module testing each comparing the patient status
18	change against an to a predetermined indicator threshold, including at least one
19	hysteresis threshold, corresponding to a same type of patient information as the
20	recorded measures which were compared[[, a]] and to at least one stickiness
21	threshold, the predetermined indicator threshold corresponding to a quantifiable
22	physiological measure of a pathophysiology indicative of angina, reduced
23	exercise capacity and respiratory distress and the hysteresis stickiness threshold
24	corresponding to a diagnosis or treatment based on a duration and degree of the
25	patient status change.
1	78. (original): A system according to Claim 77, wherein the indicator
2	thresholds relating to angina are selected from the group consisting of increased
3	ST segment decreased ST segment increased myocardial creatinine kinase.

- thresholds relating to angina are selected from the group consisting of increased ST segment, decreased ST segment, increased myocardial creatinine kinase, increased troponin, increased coronary sinus lactate, changed myocardial wall motion, ventricular tachycardia, T wave changes, and QRS changes.
- 79. (original): A system according to Claim 77, wherein the indicator thresholds relating to reduced exercise capacity are selected from the group consisting of decreased cardiac output, decreased mixed venous oxygen score, decreased patient activity score, increased pulmonary artery diastolic pressure, increased respiratory rate and decreased transthoracic impedance.

4

5

1	80. (original): A system according to Claim 77, wherein the indicator
2	thresholds relating to respiratory distress are selected from the group consisting of
3	increased pulmonary artery diastolic pressure, increased respiratory rate,
4	decreased transthoracic impedance, decreased cardiac output, decreased mixed
5	venous oxygen score, and decreased patient activity score.
1	81. (currently amended): A method for diagnosing and monitoring
2	myocardial ischemia and outcomes thereof in an automated patient care system,
3	comprising:
4	regularly recording measures relating to at least one of monitoring angina,
5	reduced exercise capacity and respiratory distress;
6	maintaining information for an individual patient, comprising:
7	organizing a plurality of monitoring sets in a database;
8	storing the recorded measures for the individual patient on a
9	substantially continuous basis into a monitoring set in the database;
10	periodically retrieving a plurality of the monitoring sets from the database
11	evaluating a finding of at least one of an absence, an onset, a progression,
12	a regression, and a status quo of myocardial ischemia, comprising:
13	determining a patient status change by comparing at least one
14	recorded measure and time from each of the monitoring sets to at least one other
15	recorded measure and time from another monitoring set with both recorded
16	measures relating to a same type of patient information recorded at different
17	times; and
18	testing each comparing the patient status change against an to a
19	predetermined indicator threshold, including at least one hysteresis threshold,
20	corresponding to a same type of patient information as the recorded measures
21	which were compared and to at least one stickiness threshold, the predetermined
22	indicator threshold corresponding to a quantifiable physiological measure of a
23	pathophysiology indicative of angina, reduced exercise capacity and respiratory

distress and the hysteresis stickiness threshold corresponding to a diagnosis or
treatment based on a duration and degree of the patient status change.
82. (original): A method according to Claim 81, wherein the indicator
thresholds relating to angina are selected from the group consisting of increased
ST segment, decreased ST segment, increased myocardial creatinine kinase,
increased troponin, increased coronary sinus lactate, changed myocardial wall
motion, ventricular tachycardia, T wave changes, and QRS changes.
83. (original): A method according to Claim 81, wherein the indicator
thresholds relating to reduced exercise capacity are selected from the group
consisting of decreased cardiac output, decreased mixed venous oxygen score,
decreased patient activity score, increased pulmonary artery diastolic pressure,
increased respiratory rate and decreased transthoracic impedance.
84. (original): A method according to Claim 81, wherein the indicator
thresholds relating to respiratory distress are selected from the group consisting of
increased pulmonary artery diastolic pressure, increased respiratory rate,
decreased transthoracic impedance, decreased cardiac output, decreased mixed
venous oxygen score, and decreased patient activity score.
85. (currently amended): A computer-readable storage medium
holding code for diagnosing and monitoring myocardial ischemia in an automated
patient care system, comprising:
code for regularly recording measures relating to at least one of
monitoring angina, reduced exercise capacity and respiratory distress;
code for maintaining information for an individual patient, comprising:
code for organizing a plurality of monitoring sets in a database;

database;

8

9

10

11

code for periodically retrieving a plurality of the monitoring sets from the

code for storing the recorded measures for the individual patient on

a substantially continuous basis into a monitoring set in the database;

code for evaluating a finding of at least one of an absence, an onset, a
progression, a regression, and a status quo of myocardial ischemia, comprising:
code for determining a patient status change by comparing at least
one recorded measure and time from each of the monitoring sets to at least one
other recorded measure and time from another monitoring set with both recorded
measures relating to a same type of patient information recorded at different
times; and
code for testing each comparing the patient status change against
an to a predetermined indicator threshold, including at least one hysteresis
threshold, corresponding to a same type of patient information as the recorded
measures which were compared and to at least one stickiness threshold, the
predetermined indicator threshold corresponding to a quantifiable physiological
measure of a pathophysiology indicative of angina, reduced exercise capacity and
respiratory distress and the hystoresis stickiness threshold corresponding to a
diagnosis or treatment based on a duration and degree of the nations status abandon